

Coupling of electrons to the electromagnetic field in a localized basis

Roland E. Allen*

Department of Physics, Texas A&M University, College Station, Texas 77843

A simple formula is obtained for coupling electrons in a complex system to the electromagnetic field. It includes the effect of intra-atomic excitations and nuclear motion, and can be applied in, e.g., first-principles-based simulations of the coupled dynamics of electrons and nuclei in materials and molecules responding to ultrashort laser pulses. Some additional aspects of nonadiabatic dynamical simulations are also discussed, including the potential of “reduced Ehrenfest” simulations for treating problems where standard Ehrenfest simulations will fail.

It is now possible to perform first-principles simulations of the coupled dynamics of electrons and nuclei with all nuclear coordinates included [1, 2, 3, 4], rather than a subset of nominal reaction coordinates. For very large systems, or when many trajectories are necessary, it is convenient to use a first-principles-based scheme [5, 6, 7, 8], with a valence-electron Hamiltonian and ion-ion repulsive potential derived from calculations using density-functional or other first-principles techniques. Here we are mainly concerned with the issue of how one can efficiently and accurately couple electrons to the electromagnetic field in such an approach, where matrix elements of various operators between localized basis functions (or “atomic orbitals”) can be calculated from first principles, and then used in large-scale calculations for complex systems, such as materials and molecules, responding to applied fields, such as ultrashort laser pulses [9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19].

Our starting point is, of course, the time-dependent Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{x}, t) = \hat{H} \psi(\mathbf{x}, t) \quad (1)$$

$$\hat{H} = \frac{1}{2m} \left(-i\hbar \nabla - \frac{q}{c} \mathbf{A}(\mathbf{x}, t) \right)^2 + U \quad , \quad q = -e. \quad (2)$$

Some time ago, Graf and Vogl [20] obtained a result, used in Refs. [13, 14, 15, 16, 17, 18, 19], which is the time-dependent version of the Peierls substitution: If \mathbf{H}_0 is the Hamiltonian matrix in a localized basis with no applied field,

$$H_0(\ell', \ell) = \int d^3x \phi_{a'}^*(\mathbf{x} - \mathbf{X}') \hat{H}_0 \phi_a(\mathbf{x} - \mathbf{X}), \quad (3)$$

and \tilde{H} is the approximate Hamiltonian when there is an applied field with vector potential $\mathbf{A}(\mathbf{x}, t)$, then they are related by

$$\tilde{H}(\ell', \ell) = H_0(\ell', \ell) e^{iq\bar{\mathbf{A}}(t) \cdot (\mathbf{X}' - \mathbf{X})/\hbar c} \quad (4)$$

with

$$\bar{\mathbf{A}}(t) = (\mathbf{A}(\mathbf{X}', t) + \mathbf{A}(\mathbf{X}, t))/2. \quad (5)$$

Here ℓ labels a localized basis function centered on a nucleus whose instantaneous position is $\mathbf{X}(\ell, t)$, and we adopt the convention of normally suppressing the indices ℓ and ℓ' , as well as the time t , by just writing \mathbf{X} and \mathbf{X}' . We will ignore any applied scalar potential A_0 , any $\boldsymbol{\mu}_B \cdot \mathbf{B}$ spin interactions, and the coupling of ion cores or nuclei to the applied fields, since these effects can be easily included when necessary.

With the prescription of (4), one does not need any new parameters in a calculation that employs either a semiempirical [13, 14] or first-principles-based [15, 16, 17, 18, 19] Hamiltonian \mathbf{H}_0 whose elements are known as a function of $(\mathbf{X} - \mathbf{X}')$. On the other hand, this prescription is in one respect a rather crude approximation: It omits intra-atomic excitations, and would therefore give no excitation at all for isolated atoms.

Here a more general version of the result of Ref. [20] will be obtained, in a form which is almost equally convenient for large-scale applications, although it does require new parameters – namely dipole matrix elements

$$\boldsymbol{\mu}_0(\ell', \ell) = q \int d^3x \phi_{a'}^*(\mathbf{x} - \mathbf{X}') (\mathbf{x} - \mathbf{X}) \phi_a(\mathbf{x} - \mathbf{X}) \quad (6)$$

and on-site ($\mathbf{X}' = \mathbf{X}$) matrix elements of the momentum operator

$$\mathbf{p}_0(\ell', \ell) = \int d^3x \phi_{a'}^*(\mathbf{x} - \mathbf{X}') (-i\hbar \nabla) \phi_a(\mathbf{x} - \mathbf{X}) \quad (7)$$

where a labels an orbital centered on the nucleus whose instantaneous position is \mathbf{X} . Recall that ℓ labels both nucleus and orbital, so at a given instant in time

$$\ell \leftrightarrow \mathbf{X}, a. \quad (8)$$

One key step is to expand ψ in terms of London orbitals, which we define to be any localized basis functions $\tilde{\phi}_a$ that are related to field-independent basis functions ϕ_a by

$$\tilde{\phi}_a(\mathbf{x} - \mathbf{X}, t) = e^{iq\mathbf{A}(\mathbf{x}, t) \cdot (\mathbf{x} - \mathbf{X})/\hbar c} \phi_a(\mathbf{x} - \mathbf{X}). \quad (9)$$

Notice that $\tilde{\phi}_a(\mathbf{x} - \mathbf{X}, t) = \phi_a(\mathbf{x} - \mathbf{X})$ when $\mathbf{A} = 0$, so that after the application of a laser pulse, for example, the London orbitals return to being standard basis functions. The ϕ_a need not be a complete set, but should, of course, be a large enough set to model all physically relevant phenomena. The relatively weak time dependence

*email allen@tamu.edu

of the nuclear positions \mathbf{X} is ignored for the moment, but will be included below. The original Hamiltonian of (2) can be rewritten as [20, 21]

$$\hat{H} = e^{iq \int \mathbf{A}(\mathbf{x}, t) \cdot d\mathbf{x}/\hbar c} \hat{H}_0 e^{-iq \int \mathbf{A}(\mathbf{x}, t) \cdot d\mathbf{x}/\hbar c} \quad (10)$$

$$\hat{H}_0 = \hat{\mathbf{p}}^2/2m + U, \quad \hat{\mathbf{p}} = -i\hbar\nabla \quad (11)$$

since (2) and (10) yield the same result when operating on an arbitrary function, and are therefore the same operator. As will be seen immediately below, there are no problems in interpreting the integral of (10) in the way that it is used here, since it is well-defined locally in evaluating each matrix element.

We now need the single approximation that $\mathbf{A}(\mathbf{x}, t)$ varies slowly with respect to \mathbf{x} over an atomic diameter or bond length, so that

$$\mathbf{A}(\mathbf{x}, t) \approx \bar{\mathbf{A}}(t) \quad (12)$$

in the matrix elements which involve $\phi_{a'}(\mathbf{x} - \mathbf{X}')$ and $\phi_a(\mathbf{x} - \mathbf{X})$. (The wavelength is thus assumed to be large compared to 1 Å.) When (10) and

$$\psi(\mathbf{x}, t) = \sum_{\ell} \psi(\ell, t) \tilde{\phi}_a(\mathbf{x} - \mathbf{X}, t) \quad (13)$$

are substituted into (1), and the resulting equation is subjected to $\int d^3x \tilde{\phi}_{a'}^*(\mathbf{x} - \mathbf{X}', t)$, we then obtain

$$\sum_{\ell} S(\ell', \ell) i\hbar \frac{\partial \psi(\ell, t)}{\partial t} = \sum_{\ell} H(\ell', \ell) \psi(\ell, t) \quad (14)$$

where

$$S(\ell', \ell) = S_0(\ell', \ell) e^{iq\bar{\mathbf{A}}(t) \cdot (\mathbf{X}' - \mathbf{X})/\hbar c} \quad (15)$$

$$S_0(\ell', \ell) = \int d^3x \tilde{\phi}_{a'}^*(\mathbf{x} - \mathbf{X}') \phi_a(\mathbf{x} - \mathbf{X}) \quad (16)$$

$$H(\ell', \ell) = \tilde{H}(\ell', \ell) - \bar{\mathbf{E}}(t) \cdot \boldsymbol{\mu}(\ell', \ell) \quad (17)$$

$$\boldsymbol{\mu}(\ell', \ell) = \boldsymbol{\mu}_0(\ell', \ell) e^{iq\bar{\mathbf{A}}(t) \cdot (\mathbf{X}' - \mathbf{X})/\hbar c} \quad (18)$$

and

$$\bar{\mathbf{E}}(t) = -\frac{1}{c} \frac{\partial \bar{\mathbf{A}}(t)}{\partial t} \quad (19)$$

is the electric field. In matrix form, (14) is

$$i\hbar \frac{\partial}{\partial t} \boldsymbol{\psi}(t) = \mathbf{S}^{-1} \cdot \mathbf{H} \cdot \boldsymbol{\psi}(t). \quad (20)$$

If there are N_e electronic basis functions, then $\boldsymbol{\psi}$ is an N_e -dimensional vector, whereas \mathbf{x} , \mathbf{A} , $\boldsymbol{\mu}$, etc. are 3-dimensional vectors. The dipole matrix elements can in principle be obtained in *ab initio* calculations like those used to obtain, e.g., the Hamiltonian matrix elements $H_0(\ell', \ell)$ [5, 6, 7, 8]. Alternatively, one might make the approximation of including only the terms with single-atom dipole matrix elements, $\boldsymbol{\mu}_0(\mathbf{X}a', \mathbf{X}a)$, and then take these from either atomic calculations or experiment.

We now return to the time dependence of the nuclear positions \mathbf{X} . With (13) rewritten as

$$\psi(\mathbf{x}, t) = \sum_{\ell} \tilde{\psi}(\ell, t) \phi_a(\mathbf{x} - \mathbf{X}) \quad (21)$$

$$\tilde{\psi}(\ell, t) = \psi(\ell, t) e^{iq\mathbf{A}(\mathbf{x}, t) \cdot (\mathbf{x} - \mathbf{X})/\hbar c} \quad (22)$$

we have [22]

$$\frac{\partial \psi}{\partial t} = \sum_{\ell} \left[\frac{\partial \tilde{\psi}(\ell)}{\partial t} \phi_a(\mathbf{x} - \mathbf{X}) + \tilde{\psi}(\ell) \frac{\partial \phi_a(\mathbf{x} - \mathbf{X})}{\partial \mathbf{X}} \cdot \dot{\mathbf{X}} \right].$$

In order to treat the second term above, we assume (as indicated by the notation) that the basis functions depend only on $(\mathbf{x} - \mathbf{X})$, so that

$$\frac{\partial \phi_a(\mathbf{x} - \mathbf{X})}{\partial \mathbf{X}} = -\frac{\partial \phi_a(\mathbf{x} - \mathbf{X})}{\partial (\mathbf{x} - \mathbf{X})} \quad (23)$$

$$= -\nabla \phi_a(\mathbf{x} - \mathbf{X}). \quad (24)$$

There is an additional correction involving $\dot{\mathbf{X}}$ which arises from

$$\begin{aligned} \frac{\partial \tilde{\psi}(\ell)}{\partial t} &= e^{iq\mathbf{A}(\mathbf{x}) \cdot (\mathbf{x} - \mathbf{X})/\hbar c} \left[\frac{\partial \psi(\ell)}{\partial t} \right. \\ &\quad \left. + \psi(\ell) \left(\frac{iq}{\hbar c} \right) \left(\frac{\partial \mathbf{A}(\mathbf{x})}{\partial t} \cdot (\mathbf{x} - \mathbf{X}) - \mathbf{A}(\mathbf{x}) \cdot \dot{\mathbf{X}} \right) \right]. \end{aligned}$$

It follows that (17) is modified to

$$\begin{aligned} H(\ell', \ell) &= H_0(\ell', \ell) e^{iq\bar{\mathbf{A}} \cdot (\mathbf{X}' - \mathbf{X})/\hbar c} \\ &\quad - \bar{\mathbf{E}}(t) \cdot \boldsymbol{\mu}(\ell', \ell) - \dot{\mathbf{X}} \cdot \mathbf{P}(\ell', \ell) \end{aligned} \quad (25)$$

where

$$\mathbf{P}(\ell', \ell) = \mathbf{p}(\ell', \ell) + (q/c) \bar{\mathbf{A}} S(\ell', \ell) \quad (26)$$

$$\mathbf{p}(\ell', \ell) = \mathbf{p}_0(\ell', \ell) e^{iq\bar{\mathbf{A}} \cdot (\mathbf{X}' - \mathbf{X})/\hbar c} \quad (27)$$

so another set of parameters is needed to treat the time dependence of the basis functions that arises from nuclear motion – namely, the matrix elements of the momentum operator $-i\hbar\nabla$.

However, when $\mathbf{X}' \neq \mathbf{X}$, there is a more convenient way of writing $\mathbf{p}_0(\ell', \ell)$:

$$\mathbf{p}_0(\ell', \ell) = i\hbar \int d^3x \tilde{\phi}_{a'}^*(\mathbf{x} - \mathbf{X}') \frac{\partial \phi_a(\mathbf{x} - \mathbf{X})}{\partial \mathbf{X}} \quad (28)$$

$$= i\hbar \frac{\partial}{\partial \mathbf{X}} S_0(\ell', \ell) \quad \text{if } \mathbf{X}' \neq \mathbf{X}. \quad (29)$$

Furthermore, in the usual case of basis functions (“atomic orbitals”) which are either even or odd under inversion through the nucleus, the fact that $(\mathbf{x} - \mathbf{X})$ and $\nabla = \partial/\partial(\mathbf{x} - \mathbf{X})$ are odd under inversion (with \mathbf{X} here taken to be fixed) implies that

$$\boldsymbol{\mu}_0(\ell, \ell) = \mathbf{p}_0(\ell, \ell) = 0. \quad (30)$$

Notice that (25) respects gauge invariance: If

$$\bar{\mathbf{A}}(t) \rightarrow \bar{\mathbf{A}}'(t) = \bar{\mathbf{A}}(t) + \Delta\bar{\mathbf{A}} \quad (31)$$

where $\Delta\bar{\mathbf{A}}$ is independent of t , then (14) still holds with

$$\psi(\ell, t) \rightarrow \psi'(\ell, t) = e^{iq\Delta\bar{\mathbf{A}} \cdot \mathbf{X}/\hbar c} \psi(\ell, t). \quad (32)$$

This is the discrete version of

$$\mathbf{A}(\mathbf{x}, t) \rightarrow \mathbf{A}'(\mathbf{x}, t) = \mathbf{A}(\mathbf{x}, t) + \nabla\Lambda(\mathbf{x}) \quad (33)$$

$$\psi(\mathbf{x}, t) \rightarrow \psi'(\mathbf{x}, t) = e^{iq\Lambda(\mathbf{x})/\hbar c} \psi(\mathbf{x}, t). \quad (34)$$

If $\Delta\bar{\mathbf{A}}$ is a function of t , gauge invariance again holds, but with the scalar potential included.

Equation (25) is the central result of the present note. This effective Hamiltonian is not manifestly Hermitian, but it still conserves probability and preserves the Pauli principle, since a straightforward calculation using (25) in (14) gives

$$i\hbar \partial \left(\psi_{n'}^\dagger \cdot \mathbf{S} \cdot \psi_n \right) / \partial t = 0 \quad (35)$$

where n labels a time-dependent one-electron state. This result also follows from the original Schrödinger equation (1) and the expansion (13), since

$$\int d^3x \psi_{n'}^*(\mathbf{x}, t) \psi_n(\mathbf{x}, t) = \psi_{n'}^\dagger(t) \cdot \mathbf{S}(t) \cdot \psi_n(t), \quad (36)$$

but it is reassuring that our approximation (12) preserves orthonormality of the time-dependent states.

For slowly moving nuclei the last term in (25) is not important. (It may be worth mentioning in this context that the direct coupling of the nuclei to the field is not considered here, since it can be treated separately.) In an earlier paper [23] we argued that the nuclear motion can be approximately treated as a “nuclear velocity field” analogous to the radiation field, and in this spirit we obtained (as a crude approximation) a generalized Peierls substitution:

$$H_{eff}(\ell', \ell) = e^{\frac{i}{\hbar} [\frac{q}{c} \mathbf{A}(\mathbf{X}') + m\dot{\mathbf{X}}'] \cdot \mathbf{X}'} \times H_0(\ell', \ell) e^{-\frac{i}{\hbar} [\frac{q}{c} \mathbf{A}(\mathbf{X}) + m\dot{\mathbf{X}}] \cdot \mathbf{X}}. \quad (37)$$

We also used this modified Hamiltonian in calculations for organic molecules responding to femtosecond-scale laser pulses of moderately strong intensity ($\sim 10^{12}$ W/cm²), and found that the $\dot{\mathbf{X}}$ terms made very little difference in the final results. On the other hand, the 2-center momentum matrix elements can be obtained from (29), and the nonzero 1-center matrix elements from either atomic calculations or experiment, so it is certainly feasible to include the last term in (25). Notice that this term is different from the Pulay correction [1], which also results from the fact that the basis functions follow the nuclei, but occurs in the equation of motion for the nuclei rather than the time-dependent Schrödinger equation for the electrons. In the kind of approach considered here

there is no Pulay correction, because the Hamiltonian matrix elements are supposed to have a position dependence that includes the movement of the basis functions.

In this context, it is worth noting that the “Ehrenfest dynamics” [24, 25] of, e.g., time-dependent density-functional theory (TDDFT) and the density-functional-based calculations of Refs. [15, 16, 17, 18, 19], can be substantially improved in molecular calculations via a trivially different procedure that might be called “reduced Ehrenfest dynamics” and which is similar in spirit to the surface hopping methods of Tully and others [26, 27]. Let us first recall some well-known results: The total wavefunction for a system of nuclei, with coordinates X_n , and electrons, with coordinates x_e , can be represented by the Born-Oppenheimer expansion

$$\Psi^{tot}(X_n, x_e, t) = \sum_i \Phi_i(X_n, t) \Psi_i(x_e, X_n). \quad (38)$$

The basis functions Ψ_i are the electronic eigenstates at fixed X_n , with the electron-nuclei and nuclei-nuclei interactions included in the electronic Hamiltonian H_e :

$$H_e(X_n) \Psi_i(x_e, X_n) = E_i(X_n) \Psi_i(x_e, X_n). \quad (39)$$

Substitution into the Schrödinger equation

$$i\hbar \partial \Psi^{tot} / \partial t = \mathcal{H} \Psi^{tot} \quad , \quad \mathcal{H} = T_n + H_e, \quad (40)$$

where T_n is the nuclear kinetic energy operator, gives an equation of the form [28, 29]

$$i\hbar \frac{\partial}{\partial t} \Phi_i = (T_n + E_i) \Phi_i - \sum_j \Lambda_{ij} \Phi_j \quad (41)$$

$$\Lambda_{ij} = \frac{\hbar^2}{2M_n} (2\mathbf{F}_{ij} \cdot \nabla_n + G_{ij}) \quad (42)$$

$$\mathbf{F}_{ij} = \langle i | \nabla_n | j \rangle \quad , \quad G_{ij} = \langle i | \nabla_n^2 | j \rangle \quad (43)$$

where M_n is a representative nuclear mass and ∇_n involves all the appropriately rescaled nuclear coordinates. If there are N_n relevant nuclear coordinates, then ∇_n and \mathbf{F} are N_n -dimensional vectors. Also, quantities in the last line are matrix elements defined in terms of Ψ_i^\dagger and Ψ_j in the usual way. If the components Φ_i are assembled into a vector Φ , (41) can be written in a form which resembles a nonabelian gauge theory [30]:

$$i\hbar \frac{\partial}{\partial t} \Phi = \left[-\frac{\hbar^2}{2M_n} (\nabla_n + \mathbf{F})^2 + \mathbf{E} \right] \cdot \Phi \quad (44)$$

where \mathbf{E} is the diagonal matrix with elements E_i . Finally, it can be shown that [31]

$$\mathbf{F}_{ij} = \frac{\langle i | \nabla_n H_e | j \rangle}{E_j - E_i} \quad , \quad E_i \neq E_j. \quad (45)$$

This last equation implies that each term in the Born-Oppenheimer expansion should evolve nearly independently if it is sufficiently distant in energy from all the other terms: If

$$|E_i - E_j| \gg |\langle i | \nabla_n H_e | j \rangle| (\hbar/P_i) \quad (46)$$

where $P_i = (2M_n E_i)^{1/2}$, then

$$i\hbar \partial \Phi_i / \partial t \approx (T_n + E_i) \Phi_i. \quad (47)$$

This is the time-dependent Born-Oppenheimer or adiabatic approximation.

On the other hand, whenever nuclear motion causes two Born-Oppenheimer “potential energy surfaces” to approach each other, so that

$$|E_i - E_j| \lesssim |\langle i | \nabla_n H_e | j \rangle| (\hbar/P_i) \quad (48)$$

there is a nonadiabatic interaction [28, 29, 32, 33], and a Born-Oppenheimer simulation based on (47) is invalid.

The results of Refs. [15, 16, 17, 18, 19] have provided a clear demonstration of the following features of simulations based on Ehrenfest dynamics:

(1) Electronic transitions are automatically observed at the points of closest approach where (48) holds, with energy released to molecular vibrations. These points are, of course, avoided crossings near the conical intersections in configuration space predicted by Teller [28, 29, 34].

(2) These transitions occur rapidly, over a time interval of ~ 1 femtosecond, during which the nuclei do not move appreciably.

Ehrenfest simulations are based on the equation of motion for the Heisenberg operator $\hat{X}(t)$ representing any nuclear coordinate [24, 25]:

$$M d^2 \hat{X} / dt^2 = -\partial \mathcal{H} / \partial \hat{X}. \quad (49)$$

Here M is the corresponding nuclear mass and \mathcal{H} is the Hamiltonian of the system. In a standard Ehrenfest simulation, the expectation value is taken over the full state of the system, including excitations (e.g. by a laser pulse) and de-excitations (e.g. by nuclear motion near conical intersections):

$$M \frac{d^2 \langle \hat{X} \rangle}{dt^2} = - \left\langle \frac{\partial \mathcal{H}(\hat{X})}{\partial \hat{X}} \right\rangle \approx - \frac{\partial \mathcal{H}(\langle \hat{X} \rangle)}{\partial \langle \hat{X} \rangle}. \quad (50)$$

There are clearly two weaknesses with this approach: First, the equality on the left represents an average over all the terms in the expansion (38), with each term representing a different nuclear trajectory. Second, the approximation on the right is totally invalid if these trajectories are very different.

Suppose, however, that the standard procedure for an Ehrenfest simulation is replaced by a trivially different procedure in which the state of the system is collapsed to a single Born-Oppenheimer term immediately after an excitation or de-excitation event. Then (47) implies that it will essentially remain in this single adiabatically evolving state until the next such event. For this reduced electronic state, the nuclei will ordinarily follow a single trajectory, except for quantum fluctuations of order $\left\langle \left(\hat{X} - \langle \hat{X} \rangle \right)^2 \right\rangle$ [35]. It is still possible for nuclear wavepackets to diverge on a single potential energy

surface, but one does not expect this to be a common occurrence for processes in which the most relevant nuclei are reasonably heavy.

For simplicity, first consider a very short laser pulse (e.g. $\sim 1 - 5$ femtoseconds in duration) applied to a molecule. The procedure for a reduced Ehrenfest simulation is as follows: Start with a single electronic eigenstate (e.g. the ground state) and initially perform an Ehrenfest simulation in the usual way. Immediately following the pulse, the molecule will be in a superposition of electronic eigenstates:

$$\Psi_e(t) = \sum_i c_i \Psi_i. \quad (51)$$

At this point one collapses Ψ_e to a single eigenstate Ψ_i and continues the simulation, with $\langle \hat{X} \rangle$ now interpreted as the expectation value for this single resulting time-dependent state, until another significant excitation or de-excitation is observed, after which there is again a further reduction to a single electronic eigenstate.

There are potentially a substantial number of branches to be followed during this sort of simulation, corresponding to the various states in the superposition (51) after an excitation or de-excitation event. The goal, however, is to understand the most relevant processes, and there will ordinarily be physical motivations for selecting the most interesting branches. Similarly, there will be many branches emerging *during* an excitation process whose duration is long enough for the nuclei to move appreciably before it is completed (e.g., a femtosecond-scale laser pulse whose duration is still $\gg 1$ femtosecond), and a choice among the branches again has to be based on physical considerations.

For a molecule subjected to high-frequency or high-intensity radiation, the branches include ionized states. The one-electron matrix element between an orbital ℓ and an ionized state with momentum \mathbf{p} is

$$H_{\ell \mathbf{p}} = \frac{e}{mc} \mathbf{A}(\mathbf{X}, t) \cdot \langle \ell | \hat{\mathbf{p}} | \mathbf{p} \rangle. \quad (52)$$

For a crude description of ionization, one might add a model orbital ϕ_0 to the basis, with

$$H_{0\ell} = \alpha_\ell \frac{e}{mc} |\mathbf{A}(\mathbf{X}, t)| p_0 \quad , \quad H_{\ell 0} = 0 \quad (53)$$

where $p_0 \sim \hbar/a_0$, a_0 is the Bohr radius, and α_ℓ is an adjustable dimensionless parameter. This non-Hermitian Hamiltonian removes amplitude from the orbital ℓ at each time step and does not return it, so it crudely models excitation to a localized wavepacket with the electron ultimately escaping the system. An appreciable probability for a given ionized state then provides motivation for following that branch in a reduced Ehrenfest simulation. Notice that an accurate treatment of ionization is not necessary if the only issue is whether an ionized state is important enough to warrant a simulation of the subsequent dynamics in that state. Also notice that the

energy H_{00} of the extra orbital is irrelevant (so one can take $H_{00} = 0$) and that a single extra orbital is sufficient regardless of the size of the system.

After each wavefunction collapse, the use of (50) implies that the nuclei are treated classically. It is then appropriate to use the mixed classical-quantum action [25, 36] $S = \int dt L$, where

$$L = \frac{1}{2} \langle \Psi_e | \left(i\hbar \frac{\partial}{\partial t} - \mathcal{H}_e \right) | \Psi_e \rangle + h.c. + \frac{1}{2} \sum_{k\alpha} M_k \left(\frac{dX_{k\alpha}}{dt} \right)^2 - U_{rep} \quad (54)$$

where \mathcal{H}_e is the electronic Hamiltonian, $|\Psi_e\rangle$ is the electronic state, “h.c.” means “Hermitian conjugate”, k labels a nucleus with spatial coordinates α , and U_{rep} is the repulsive interaction between nuclei or ion cores. As shown in Ref. [36] (but with \mathbf{H} now given by (25)), extremalization of this action leads to (20) and

$$M \frac{d^2 X}{dt^2} = -\frac{1}{2} \sum_n \psi_n^\dagger \cdot \left(\frac{\partial \mathbf{H}}{\partial X} - i\hbar \frac{\partial \mathbf{S}}{\partial X} \frac{\partial}{\partial t} \right) \cdot \psi_n + h.c. - \frac{\partial U_{rep}}{\partial X} \quad (55)$$

if one makes the usual time-dependent effective-field approximation, with exchange and correlation represented by an effective one-electron potential, and the electronic state represented by a single antisymmetrized product wavefunction $\Psi_e(t)$. Here X is any nuclear coordinate and M is the corresponding mass.

The reduced Ehrenfest method described above combines the advantages of Born-Oppenheimer simulations, which are valid when (47) holds, and Ehrenfest simulations, which are suitable for treating the vibronic transitions when (48) holds, as the results of Refs. [15, 16, 17, 18, 19] have clearly demonstrated. The use of reduced

Ehrenfest simulations should solve various problems that are encountered in standard Ehrenfest simulations – for example, the apparent failure of TDDFT to correctly describe the isomerization of retinal [8]. One problem with TDDFT is that the energies of excited states are not accurately described, but a potentially more severe problem in the case of molecules is that TDDFT is a special case of standard Ehrenfest dynamics, and as a result fails to yield a complete return to the ground state following de-excitation near a conical intersection. In a reduced Ehrenfest simulation, on the other hand, one correctly follows the nuclear dynamics for that fraction of the population of molecules which does return to the ground state, and which therefore should isomerize more readily. Reduced Ehrenfest simulations are practical for large molecules, and are still consistent with the true meaning of quantum amplitudes, which yield probabilities for the various outcomes that are observed at the classical level.

Finally, it may be worth noting that the above treatment can be straightforwardly generalized to other particles, relativistic systems, and nonabelian gauge fields, with ψ in (1) interpreted as a multicomponent field and the Hamiltonian of (2) appropriately changed. It can also be used with many-body effects included through self-energy terms, in the Kadanoff-Baym/Keldysh equations for time-dependent and nonequilibrium problems [37, 38]. The chief limitation is the use of localized basis functions and the approximation (12).

Acknowledgements

This work was supported by the Robert A. Welch Foundation (Grant A-0929). I also thank Yusheng Dou and Meng Gao for their helpful comments.

-
- [1] M. C. Payne, M. P. Teter, D. C. Allan, T. A. Arias, and J. D. Joannopoulos, *Rev. Mod. Phys.* 64, 1045 (1992).
 - [2] J. Li, G. Speyer, and O. F. Sankey, *Phys. Rev. Lett.* 93, 248302 (2004).
 - [3] J. Van Heys, M. Lindenblatt, and E. Pehlke, *Phase Transitions* 78, 773 (2005).
 - [4] *Time-Dependent Density Functional Theory*, edited by M. A. L. Marques, C. A. Ullrich, F. Nogueira, A. Rubio, K. Burke, and E. K. U. Gross (Springer, Heidelberg, 2006).
 - [5] D. Porezag, Th. Frauenheim, Th. Köhler, G. Seifert, and R. Kaschner, *Phys. Rev. B* 51, 12947 (1995).
 - [6] M. Elstner, D. Porezag, G. Jungnickel, J. Elsner, M. Haugk, Th. Frauenheim, S. Suhai, and G. Seifert, *Phys. Rev. B* 58, 7260 (1998).
 - [7] Th. Frauenheim, G. Seifert, M. Elstner, T. Niehaus, C. Köhler, M. Amkreutz, M. Sternberg, Z. Hajnal, A. D. Carlo, and S. Suhai, *J. Phys. Condens. Matt.* 14, 3015 (2002).
 - [8] T. A. Niehaus, D. Heringer, B. Torralva, and Th. Frauenheim, *Eur. Phys. J. D* 35, 467 (2005).
 - [9] S. K. Sundaram and E. Mazur, *Nature Materials* 1, 217 (2002).
 - [10] A. M. Lindenberg, J. Larsson, K. Sokolowski-Tinten, K. J. Gaffney, C. Blome, O. Synnergren, J. Sheppard, C. Caleman, A. G. MacPhee, D. Weinstein, D. P. Lowney, T. K. Allison, T. Matthews, R. W. Falcone, A. L. Cavalieri, D. M. Fritz, S. H. Lee, P. H. Bucksbaum, D. A. Reis, J. Rudati, P. H. Fuoss, C. C. Kao, D. P. Siddons, R. Pahl, J. Als-Nielsen, S. Duesterer, R. Ischebeck, H. Schlarb, H. Schulte-Schrepping, Th. Tschentscher, J. Schneider, D. von der Linde, O. Hignette, F. Sette, H. N. Chapman, R. W. Lee, T. N. Hansen, S. Techert, J. S. Wark, M. Bergh, G. Huld, D. van der Spoel, N. Timneanu, J. Hajdu, R. A. Akre, E. Bong, P. Krejčík, J. Arthur, S. Brennan, K. Luening, J. B. Hastings, *Science* 308, 392 (2005).

- [11] P. B. Hillyard, K. J. Gaffney, A. M. Lindenberg, S. Engemann, R. A. Akre, J. Arthur, C. Blome, P. H. Bucksbaum, A. L. Cavalieri, A. Deb, R. W. Falcone, D. M. Fritz, P. H. Fuoss, J. Hajdu, P. Krejcik, J. Larsson, S. H. Lee, D. A. Meyer, A. J. Nelson, R. Pahl, D. A. Reis, J. Rudati, D. P. Siddons, K. Sokolowski-Tinten, D. von der Linde, and J. B. Hastings, *Phys. Rev. Lett.* **98**, 125501 (2007).
- [12] A. H. Zewail, *Femtochemistry: Ultrafast Dynamics of the Chemical Bond* (World Scientific, Singapore, 1994); *Angew. Chem. Int. Ed.* **39**, 2586 (2000).
- [13] J. S. Graves and R. E. Allen, *Phys. Rev. B* **58**, 13627 (1998).
- [14] T. Dumitrica and R. E. Allen, *Phys. Rev. B* **66**, 081202(R) (2002).
- [15] B. Torralva, T. A. Niehaus, M. Elstner, S. Suhai, Th. Frauenheim, and R. E. Allen, *Phys. Rev. B* **64**, 153105 (2001).
- [16] Y. Dou and R. E. Allen, *Chem. Phys. Lett.* **378**, 323 (2003).
- [17] Y. Dou, B. R. Torralva, and R. E. Allen, *Chem. Phys. Lett.* **392**, 352 (2004).
- [18] P. Sauer and R. E. Allen, *Chem. Phys. Lett.* **434**, 260 (2007).
- [19] P. Sauer and R. E. Allen, *Chem. Phys. Lett.* **450**, 192 (2008).
- [20] M. Graf and P. Vogl, *Phys. Rev. B* **51**, 4940 (1995).
- [21] R. Peierls, *Z. Phys.* **80**, 763 (1933).
- [22] T. N. Todorov, *J. Phys. Condens. Matt.* **13**, 10125 (2001).
- [23] T. Dumitrica, A. Burzo, Y. Dou, and R. E. Allen, *Phys. Stat. Sol. (b)* **241**, 2331 (2004).
- [24] J. J. Sakurai, *Modern Quantum Mechanics* (Addison-Wesley, Reading, Massachusetts, 1994).
- [25] R. E. Allen, *Phys. Rev. B* **50**, 18629 (1994).
- [26] J. C. Tully, *J. Chem. Phys.* **93**, 1061 (1990).
- [27] G. Granucci and M. Persico, *J. Chem. Phys.* **126**, 134114 (2007) and references therein.
- [28] *Conical Intersections: Electronic Structure, Dynamics, and Spectroscopy*, edited by W. Domcke, D. R. Yarkony, and H. Köppel (World Scientific, Singapore, 2004).
- [29] M. Baer, *Beyond Born-Oppenheimer: Electronic Nonadiabatic Coupling Terms and Conical Intersections* (Wiley Interscience, Hoboken, 2006).
- [30] B. Zygelman, *Phys. Lett. A* **125**, 476 (1987).
- [31] S. T. Epstein, *Am. J. Phys.* **22**, 613 (1954).
- [32] L. S. Cederbaum, *J. Chem. Phys.* **128**, 124101 (2008) and references therein.
- [33] M. Ben-Nun, J. Quenneville, and T. J. Martínez, *J. Phys. Chem. A* **104**, 5161 (2000).
- [34] E. Teller, *J. Phys. Chem.* **41**, 109 (1937).
- [35] P. Sauer, Y. Rostovtsev, and R. E. Allen, *J. Chem. Phys.* **126**, 024502 (2007).
- [36] R. E. Allen, T. Dumitrica, and B. Torralva, Chapter 7 of *Ultrafast Physical Processes in Semiconductors*, edited by K. T. Tsen (Academic, New York, 2001).
- [37] *Introduction to Computational Methods in Many Body Physics*, edited by M. Bonitz and D. Semkat (Rinton Press, Paramus, New Jersey, 2006) and references therein.
- [38] N. E. Dahlen and R. van Leeuwen, *Phys. Rev. Lett.* **98**, 153004 (2007).